



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

# Management Of Experiments And Data At The National Ignition Facility

S. Azevedo, A. Casey, R. Beeler, R. Bettenhausen, E. Bond, H. Chandrasekaran, C. Foxworthy, M. Hutton, J. Krammen, J. Liebman, A. Marsh, T. Pannell, J. Rhodes, J. Tapperro, A. Warrick

March 25, 2011

ICALEPCS  
Grenoble, France  
October 10, 2011 through October 14, 2011

## **Disclaimer**

---

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

# **MANAGEMENT OF EXPERIMENTS AND DATA AT THE NATIONAL IGNITION FACILITY\***

S. Azevedo, A. Casey, R. Beeler, R. Bettenhausen, E. Bond, H. Chandrasekaran, C. Foxworthy, M. Hutton, J. Krammen, J. Liebman, A. Marsh, T. Pannell, J. Rhodes, J. Tapper, A. Warrick

Lawrence Livermore National Laboratory, P.O. Box 808, Livermore, California USA

## **ABSTRACT**

Experiments, or “shots”, conducted at the National Ignition Facility (NIF) are discrete events that occur over a very short time frame (tens of nanoseconds) separated by many hours. Each shot is part of a larger campaign of shots to advance scientific understanding in high-energy-density physics. In one campaign, scientists use energy from the 192-beam, 1.8-Megajoule pulsed laser in the NIF system to symmetrically implode a hydrogen-filled target, thereby creating conditions similar to the interior of stars in a demonstration of controlled fusion. Each NIF shot generates gigabytes of data from over 30 diagnostics that measure optical, x-ray, and nuclear phenomena from the imploding target. We have developed systems to manage all aspects of the shot cycle. Other papers will discuss the control of the lasers and targets, while this paper focuses on the setup and management of campaigns and diagnostics. Because of the low duty cycle of shots, and the thousands of adjustments for each shot (target type, composition, shape; laser beams used, their power profiles, pointing; diagnostic systems used, their configuration, calibration, settings) it is imperative that we accurately define all equipment prior to the shot. Following the shot, and capture of the data by the automatic control system, it is equally imperative that we archive, analyze and visualize the results within the required 30 minutes post-shot. Results must be securely archived, approved, web-visible and downloadable in order to facilitate subsequent publication. To-date NIF has successfully fired over 2,500 system shots, as well as thousands of test firings and dry-runs. We will present an overview of the highly-flexible and scalable campaign management systems and tools employed at NIF that control experiment configuration of the facility all the way through presentation of analyzed results.

\* This work performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

284 words, 1597 characters